arXiv:1205.6300v2 [math.PR] 15 Oct 2012

Extremal Lipschitz functions in the deviation inequalities from the mean

Dainius Dzindzalieta[1]

March 5, 2013

Abstract

We consider a deviation from the mean in the problem

$$D(x) \overset{\mathrm{def}}{=} \sup_{f\in\mathcal{F}} \mu\{f - \mathbf{E}_\mu f \geq x\}, \qquad \text{for } x \in \mathbf{R} \tag{0.1}$$

for the class $\mathcal{F}$ of all integrable, $f \in L_1(V, d, \mu)$, Lipschitz functions on a probability metric space (V, d, μ). Consider the following closely related (vertex-)isoperimetric problem. Given $t > 0$,

$$\text{minimize } \mu(A^h) \text{ over all } A \subset V \text{ with } \mu(A) \geq t, \tag{0.2}$$

where $A^h = \{u \in V : d(u, A) \leq h\}$ is an h-neighborhood of A. We solve the problem (0.1) for spaces (V, d, μ) such that for every $t \geq 0$ there exists a solution of (0.2) which does not depend on h. As corollaries we get exact solutions of (0.1) for Euclidean unit sphere S^{n-1} with a geodesic distance and a normalized Haar measure, for $\mathbf{R}^n$ equipped with a Gaussian measure and for n-dimensional cube, rectangle, powers of tori, Diamond graph equipped with uniform measure and Hamming distance. We also obtain a general result for probability metric spaces. Let $\mathcal{F}^*$ be a family of distance functions $f(u) = -d(A, u)$, with $A \subset V$ measurable. For all probability metric spaces we prove that

$$\sup_{f\in\mathcal{F}} \mu\{f - \mathbf{E}_\mu f \geq x\} = \sup_{f\in\mathcal{F}^*} \mu\{f - \mathbf{E}_\mu f \geq x\}, \qquad x \in \mathbf{R}.$$

2010 *Mathematics Subject Classification*: Primary 60E15; Secondary 60A10.

[1] Institute of Mathematics and Informatics, Vilnius University, Vilnius, Lithuania. E-mail: dainiusda@gmail.com

[1] This research was funded by a grant (No. MIP-12090) from the Research Council of Lithuania

Key words and phrases: Gaussian, vertex isoperimetric, deviation from the mean, inequalities, Hamming, probability metric space.

1 Introduction and results

For a metric space (V,d), we denote by $\mathcal{F} = \mathcal{F}(V)$ the class of integrable, i.e. $f \in L_1(V,d,\mu)$, 1-Lipschitz functions $f : V \to \mathbf{R}$, i.e., such that $|f(u) - f(v)| \le d(u,v)$ for all $u,v \in V$. In this paper we consider a deviation from the mean in the problem

$$D(x) \overset{\text{def}}{=} \sup_{f \in \mathcal{F}} \mu\{f - \mathbf{E}_\mu f \ge x\}, \qquad \text{for } x \in \mathbf{R} \tag{1.1}$$

for the class $\mathcal{F}$ of all Lipschitz functions on a probability metric space (V,d,μ). Here a probability metric space means that a measure μ is Borel and normalized, $\mu(V) = 1$. If we change f to $-f$ we get that

$$D(x) = \sup_{f \in \mathcal{F}} \mu\{f - \mathbf{E}_\mu f \le -x\} \qquad \text{for } x \in \mathbf{R}.$$

Note that the function $D(x)$ depends also on (V,d,μ).

In order to state our results we need some notation and preliminaries. Consider the following (vertex-)isoperimetric problem. Given t,

$$\text{minimize } \mu(A^h) \text{ over all } A \subset V \text{ with } \mu(A) \ge t, \tag{1.2}$$

where $A^h = \{u \in V : d(u,A) \le h\}$ is an h-extension of A. A nice overview of isoperimetric problems with applications can be found in [Led01, LT91] e.g. Following [BS02] we say that a space (V,d,μ) is *isoperimetric* if for every $t \ge 0$ there exists a solution, say A_{opt}, of (1.2) which does not depend on h. We derive exact deviation from the mean inequalities for Lipschitz functions on isoperimetric probability metric spaces.

We firstly state a general result for all probability metric spaces. It states that, the sup in (1.1) is achieved on a family, say $\mathcal{F}^*$, of distance functions $f(u) = -d(A,u)$ with measurable $A \subset V$. Note that $\mathcal{F}^* \subset \mathcal{F}$.

Theorem 1.1. *Given a probability metric space (V,d,μ), we have*

$$\sup_{f \in \mathcal{F}} \mu\{f - \mathbf{E}_\mu f \ge x\} = \sup_{f \in \mathcal{F}^*} \mu\{f - \mathbf{E}_\mu f \ge x\} \qquad x \in \mathbf{R}.$$

Proof Fix $x \in \mathbf{R}$. Let $f \in \mathcal{F}$ and $B = \{u : f(u) - \mathbf{E}_\mu f \ge x\}$. Without loss of generality we suppose that $f \ge 0$ for $u \in B$ and $f(u) = 0$ for at least one $u \in B$. Let $g(u) = -d(B,u)$. It is easy to see that $f(u) \ge g(u)$ for all $u \in V$ and $\mathbf{E}_\mu f \ge \mathbf{E}_\mu g$, so $B \subset \{u : g(u) - \mathbf{E}_\mu g \ge x\}$. Thus $\mu\{f - \mathbf{E}_\mu f \ge x\} \le \mu\{g - \mathbf{E}_\mu g \ge x\}$. Since f is arbitrary the statement of *Theorem* 1.1 follows. ∎

In the special case when $V = \mathbf{R}^n$ and $\mu = \gamma_n$ is a standard Gaussian measure, $Theorem$ 1.1 was proved by Bobkov [Bob03].

Using $Theorem$ 1.1 we solve the problem (1.1) for all isoperimetric probability metric spaces.

Theorem 1.2. *For all isoperimetric probability metric spaces* (V, d, μ) *we have*

$$D(x) = \mu\{f^* - \mathbf{E}_\mu f^* \geq x\} \qquad \text{for } x \in \mathbf{R},$$

where $f^*(u) = d(A_{opt}, u)$ *is a distance function from some extremal set* A_{opt}*. It turns out that* $\mu(A_{opt}) = D(x)$*.*

Proof We have

$$\begin{aligned} \mathbf{E}_\mu d(A, \cdot) &= \int_0^\infty \left(1 - \mu\left\{A^h\right\}\right) \mathrm{dh} \\ &\leq \int_0^\infty \left(1 - \mu\left\{A^h_{\mathrm{opt}}\right\}\right) \mathrm{dh} = \mathbf{E}_\mu d(A_{\mathrm{opt}}, \cdot). \end{aligned}$$

Thus by $Theorem$ 1.1

$$\sup_{f \in \mathcal{F}} \mu\{f - \mathbf{E}_\mu f \geq x\} = \sup_{f \in \mathcal{F}^*} \mu\{f - \mathbf{E}_\mu f \geq x\} \leq \mu\{f^* - \mathbf{E}_\mu f^* \geq x\}$$

with $f^*(u) = -d(A_{\mathrm{opt}}, u)$. Which completes the proof of $Theorem$ 1.2. ∎

2 Isoperimetric spaces and corollaries

In this section we provide a short overview of the most known and important isoperimetric spaces. We also state a number of corollaries following from $Theorem$ 1.1 and $Theorem$ 1.2.

A typical and basic example of isoperimetric spaces is the Euclidean unit sphere $S^{n-1} = \{x \in \mathbf{R}^n : \sum_{i=1}^n |x_i|^2 = 1\}$ with a geodesic distance ρ and a normalized Haar measure σ_{n-1}. P. Lévy [Lév51] and E. Schmidt [Sch48] showed that if A is a Borel set in S^{n-1} and H is a cap (ball for geodesic distance ρ) with the same Haar measure $\sigma_{n-1}(H) = \sigma_{n-1}(A)$, then

$$\sigma_{n-1}(A^h) \geq \sigma_{n-1}(H^h) \qquad \text{for all } r > 0. \tag{2.1}$$

Thus A_{opt} for the space $(S^{n-1}, \rho, \sigma_{n-1})$ is a cap. We refer readers for short proof of (2.1) to [FLM77, Ben84]. See the extension to Riemannian manifolds with strictly positive curvature in [Gro80]. Note that if H is a cap, then H^h is also a cap, so we have an immediate corollary.

Corollary 2.1. *For a unit sphere S^{n-1} with geodesic distance equipped with normalized Haar measure σ_{n-1} we have*

$$D(x) = \sigma_{n-1}\{f^* - \mathbf{E}_\mu f^* \geq x\} \qquad \text{for } x \in \mathbf{R},$$

where $f^*(u) = -d(A_{opt}, u)$ and A_{opt} is a cap.

Probably the most simple non-trivial isoperimetric space is n-dimensional discrete cube $C_n = \{0,1\}^n$ with usual Hamming distance equipped with uniform measure, say μ. Since μ is uniform we have $\mu\{u\} = 1/2^n$ for all $u \in C_n$. Harper [Har64] proved that the set of some first elements of C_n in lexicographical order is a solution of (1.2). Other well studied discrete isoperimetric spaces with uniform measures are powers of tori [Kar82, Rio98], n-dimensional rectangle [BL91], powers of the Diamond graph [BRS08] we also mention powers of so called cross-sections [BS02]. We state the results for discrete spaces as corollary.

Corollary 2.2. *For discrete n-dimensional cube, n-dimensional rectangle, powers of tori and Diamond graphs with usual Hamming distance equipped with uniform measure we have*

$$D(x) = \mu\{f^* - \mathbf{E}_\mu f^* \geq x\} \qquad \text{for } x \in \mathbf{R},$$

where $f^*(u) = -d(A_{opt}, u)$ and A_{opt} are the sets of some first elements in corresponding orders. In particular, for n-dimensional discrete cube with Hamming distance A_{opt} is a set of some first elements of C_n in lexicographical order.

There is a vast of papers dedicated to bound $D(x)$ for various discrete spaces. We mention only [BH97, Led01, LT91, Tal95] among others. In [LT91, BRS08] a nice overview of isoperimetric spaces and bounds for $D(x)$ are provided.

Another important example of isoperimetric spaces comes from Gaussian isoperimetric problem. Sudakov and Tsirel'son [ST78] and Borell [Bor75] discovered that if γ_n is a standard Gaussian measure on $\mathbf{R}^n$ with a usual Euclidean distance function d, then $(\mathbf{R}^n, d, \gamma_n)$ is isoperimetric. In [ST78] and [Bor75] it was shown that among all subsets A of $\mathbf{R}^n$ with $t \geq \gamma_n(A)$, the minimal value of $\gamma_n(A^h)$ is attained for half-spaces of measure t. Thus we have the following corollary of *Theorem* 1.1 and *Theorem* 1.2.

Corollary 2.3. *For a Gaussian space* $(\mathbf{R}^n, d, \gamma_n)$ *we have*

$$D(x) = \gamma_n\{f^* - \mathbf{E}_{\gamma_n} f^* \geq x\} \qquad \textit{for } x \in \mathbf{R},$$

where $f^*(u) = -d(A_{opt}, u)$ *is a distance function from a half-space of space* $\mathbf{R}^n$.

This corollary was firstly proved by Bobkov [Bob03]. We also refer for further investigations of extremal sets on $\mathbf{R}^n$ for some classes of measures to [Bob97, BCR07] among others.

References

[BCR07] F. Barthe, P. Cattiaux, and C. Roberto. Isoperimetry between exponential and Gaussian. *Electron. J. Probab*, 12(44):1212–1237, 2007.

[Ben84] Y. Benyamini. Two-point symmetrization, the isoperimetric inequality on the sphere and some applications, in Texas Functional Analysis Seminar 1983/84, Austin, Texas, Longhorn Notes, 1984.

[BH97] S.G. Bobkov and C. Houdré. Isoperimetric constants for product probability measures. *The Annals of Probability*, 25(1):184–205, 1997.

[BL91] B. Bollobás and I. Leader. Isoperimetric inequalities and fractional set systems. *Journal of Combinatorial Theory, Series A*, 56(1):63–74, 1991.

[Bob97] S.G. Bobkov. Isoperimetric problem for uniform enlargement. *Studia Math*, 123(1):81–95, 1997.

[Bob03] S.G. Bobkov. Localization Proof of the Bakry–Ledoux Isoperimetric Inequality and Some Applications. *Theory of Probability and its Applications*, 47:308, 2003.

[Bor75] C. Borell. The Brunn-Minkowski inequality in gauss space. *Inventiones Mathematicae*, 30(2):207–216, 1975.

[BRS08] S.L. Bezrukov, M. Rius, and O. Serra. The vertex isoperimetric problem for the powers of the diamond graph. *Discrete Mathematics*, 308(11):2067–2074, 2008.

[BS02] S.L. Bezrukov and O. Serra. A local-global principle for vertex-isoperimetric problems. *Discrete Mathematics*, 257(2):285–310, 2002.

[FLM77] T. Figiel, J. Lindenstrauss, and VD Milman. The dimension of almost spherical sections of convex bodies. *Acta Mathematica*, 139(1):53–94, 1977.

[Gro80] M. Gromov. Paul Levy's isoperimetric inequality. *Preprint I.H.E.S*, 1980.

[Har64] L.H. Harper. Optimal assignments of numbers to vertices. *Journal of the Society for Industrial and Applied Mathematics*, 12(1):131–135, 1964.

[Kar82] V.M. Karachanjan. A discrete isoperimetric problem on multidimensional torus. In *Russian), Doklady AN Arm. SSR, vol. LXXIV*, volume 2, pages 61–65, 1982.

[Led01] M. Ledoux. *The concentration of measure phenomenon.* Amer Mathematical Society, 2001.

[Lév51] Paul Lévy. *Problèmes concrets d'analyse fonctionnelle. Avec un complément sur les fonctionnelles analytiques par F. Pellegrino.* Gauthier-Villars, Paris, 1951. 2d ed.

[LT91] M. Ledoux and M. Talagrand. *Probability in Banach Spaces: isoperimetry and processes.* Springer, 1991.

[Rio98] O. Riordan. An ordering on the even discrete torus. *SIAM Journal on Discrete Mathematics*, 11:110, 1998.

[Sch48] E. Schmidt. Die Brunn-Minkowskische Ungleichung und ihr Spiegelbild sowie die isoperimetrische Eigenschaft der Kugel in der euklidischen und nichteuklidischen Geometrie. I. *Mathematische Nachrichten*, 1(2-3):81–157, 1948.

[ST78] V.N. Sudakov and B.S. Tsirel'son. Extremal properties of half-spaces for spherically invariant measures. *Journal of Mathematical Sciences*, 9(1):9–18, 1978.

[Tal95] M. Talagrand. Concentration of measure and isoperimetric inequalities in product spaces. *Publications Mathematiques de l'IHES*, 81(1):73–205, 1995.